



EFFECT OF ANTS GENERA ON SUSTAINING MODIFICATION AND FERTILITY OF SOIL

SEEMA. G. KADU

Assistant Prof. Dept. of Zoology,
S.S.E.S. Shri Shivaji Science College, Congress Nagar Nagpur,
seema.kadu75@gmail.com

Abstract

Ants are important in below ground processes through the alteration of the physical and chemical environment and through their effects on plants, microorganisms, and other soil organisms. The role of ants in ecosystems is discussed, mainly from the perspective of the effects of ground-dwelling ants on soil processes and function, emphasizing their role as ecosystem engineers. Different ant species navigate across all the trophic levels in search of architecture of vegetation, dead twigs on soil for nesting. An ant represents a unique focal group having the potential to be used as biological indicators. The maximum population of ant species effects on the physical and biological properties of soil greatly. Also, the movement of particles from lower horizons to the surface area by ants aids affects the mixing of organic and mineral fractions of the soil to increase the fertility. The faeces of ants are the basis for the formation of soil aggregates and humus, which physically stabilize the soil and increase its porosity to store nutrients. The study of importance of ants in ecosystem functioning, principally their ability to maintain or restore soil quality as bio-indicators. Present study examined the mound soil of ant species acting as prodigious and legendary soil bioturbator involved in intensification in concentration of soil nutrients N, P(PO⁴), S(SO⁴⁺), Mg²⁺, Ca²⁺, and K⁺.

Keywords: Ant, porosity, nutrients, fertility Bioturbator.

Introduction

Some ants like weaver ants, pharaoh ants, wood cutter ant and carpenter ant species construct their nest in the soil to deposit their food and egg laying playing an important role in the basic nutrient cycle of the soil. As plants grow, they consume nutrients in the soil. As the mineral content depletes in the soil, they need to be replenished. This is where the organisms living in the soil play their part by helping in the decomposition and transportation of nutrients to be used by plants. As some wood ants are keystone species in woodland ecosystems, with effects on the community structure of local invertebrates as well as providing a food source for predators (Hughes and Broome, 2007). Nest construction results in modification of soil structure, increasing porosity (Frouz and Jilková, 2008) and accumulation of food and detritus makes nests hotspots of nutrient exchange (Domisch et al., 2009). The nests constructed by these ants support high levels of biodiversity, including many species that are dependent on the nests as habitat (Wagner and Jones 2005). There are different ant species shows burrowing activity among these the ants species under subfamilies Dolichoderinae, Myrmicinae, Pseudomyrmicinae, Ponerinae, Formicinae, are the important in soil tilting during nest building. These ants genera plays important role to create

porous soil, provide better aeration, improves the moisture content and helps in the transfer of organic materials into the soil. Among these species some genera, *Iridomyrmex*, *Tapinoma*, *Camponotus*, *Oecophylla*, *Paratrechina*, *Aphaenogaster*, *Crematogaster*, *Solenopsis*, *Pachycondyla*, are the polymorphic hymenopterans utilizes the plant residues, animal manure or other organic materials to build up their nest colony (Frouz *et.al.*, 2003). These organic materials decomposed by bacteria, fungi and converts the complex organic matter into simple components (Eldridge and Pickard, 1994). The population of these ants genera are directly effects on the physical, chemical and biological properties of soil through the modification of soil profile, aeration, drainage and density and are therefore able to influence the energy and nutrient cycles and to act as ecological engineers (Folgarait, 1998; Frouz and Jilkova, 2008). The reduced simple organic components are again carried by the ants in to excavated nest underground the soil where they are further being absorbed by the plants roots as growth nutrients (Cammeraat *et.al.*, 2002).

Material and Method

This experiment was conducted to find out whether the fauna inhabiting the soil will help with soil fertility and plant growth. The experiment was carried out in some rural villages near Kalmeshwar region in Nagpur District (MS). The five different experimental areas near the vegetation owning maximum number of ant mound of different ant genera under subfamilies Dolichoderinae, Myrmicinae, Pseudomyrmicinae, Ponerinae and Formicinae are studied as the experimental plot. The soil sample collected from roof of the plant and tested in the laboratory in college using Soil Testing Kit. The phosphorus concentration in soil tests measure ortho-phosphate (PO₄-P) and ammonium acetate extraction method is the most common method to analyse potassium concentration in ppm. The vegetative areas without ant mound and ant population were studied as check or control.

Results and Discussion

The ant genera under subfamilies Dolichoderinae, Myrmicinae, Pseudomyrmicinae, Ponerinae, Formicinae, play very important role in soil tilting during nest building. Also the present study focused on the important role of ants to create porous soil, provide better aeration, improves the moisture content and helps in the transfer of organic materials into the soil. It is observed that the soil collected from different areas denoted as A-1, A-2, A-3, A-4 and A-5 had grown taller than the control area denoted as A-c. The presence of these ants genera in areas A-1 to A-5 show increasing height and growth of plant population. The hypothesis that the presence of these ants in the soil help to provide a better environment for the plant growth. The concentration of Potassium (K), Phosphorous (P) and pH in the respective soil sample from A-1 to A-5 areas show variable values as compared to control area (Table-1).

Table-1 : Concentration of Soil Nutrients (in ppm) at experimental ant mound areas (A=05).

S. No.	Concentration of Soil Nutrients (ppm)	A-1	A-2	A-3	A-4	A-5	Control A-C
1.	N	55>	65>	70>	42>	58>	30 <
2.	P(PO₄)	54>	68>	72>	62>	55>	31<
3.	S(SO₄²⁻)	20.5>	25>	20>	22>	25>	11.5<
4.	Mg²⁺	188>	155>	167>	175>	180>	70 <
5.	Ca²⁺	2300>	2200>	1800>	2100>	2300>	1200<
6.	K⁺	140>	165>	186>	195>	202>	105<

P<0.001

Conclusion:

Some of the ants eat other insects and caterpillars that are harmful to the plants and will help nourish the soil with their processed food (Barsagade et al., 2013). The burrowing activity of the ant genera helps to penetrate water up to the plants roots (Domisch, 2006). The present study shows burrowing habitat and mining activity among some ant genera under subfamilies Dolichoderinae, Myrmicinae, Pseudomyrmicinae, Ponerinae, Formicinae, are very important to play an important role to create porous soil, provide better aeration, improve the moisture content and help in the transfer of organic materials into the soil to fertilize the soil in many ways (Frouz et al., 2003). They help in the recycling of decomposed materials which increase soil porosity and may cause separation of soil particles according to their size. The ants also help in spreading the plants seeds during these soil mining. Ant-mediated chemical changes of soil are represented mainly by a shift of pH towards neutral and an increase in nutrient content (mostly Potassium and phosphorus) in ant nest-affected soil (Frouz and Jilkova, 2008). Effects on the soil vary between ant species; substantial variation can be found in the same species living in different conditions. The increase in concentration of Potassium and phosphorus and neutral pH represent an ideal indicator for experiments targeting soil ecosystems (Lafleur, 2005). These effects correspond with accumulation of food in the nests and the effect on biological processes, such as hastening of decomposition rate (Kadu, 2010, 2014). Effects on biological soil properties may be connected with increased or decreased microbial activity, which is affected by accumulation of organic matter and internal nest temperature and especially moisture (Eldridge and Myers 1998). In this study, we compared nutrient exploitation by spruce seedlings growing in substrates from abandoned ant nests (i.e., local nutrient-rich hot-spots) vs. from the control soil floor area (Ohashi et al., 2007). At field moisture, microbial biomass Carbon and Nitrogen were significantly more concentrated in ant nests at all experimental study sites (A-1 to A-5) as compared to control sites (Frouz et al., 2008). The soil becomes naturally moistened in the form of nest soils and had higher capacity for microbial growth than soils from other microhabitats (Wagner and Jones, 2005). However, this study is the first to document the direct linkage of select nutrient availability in ant mound soils as well as soil bioturbation with enhanced nutrient levels expressed in perimeter vegetation. This concludes that ant nest mounds highly impact on the soil tilting by creating concentrated patches of soil with increase in nutrients concentration could affect biogeochemical cycling rates and plant community dynamics.

Acknowledgement

Author greatly thankful to Dr. D.B. Tembhare , Ex. Prof. and Head of Department of Zoology, Campus and Dr. D. D. Barsagade, Reader, Department of Zoology, RTM Nagpur Univ. Nagpur.

Reference:-

Avinoam Danin and Yom Tov. 1990. Ant nests as primary habitats of *Silybum marianum* (Compositae). *Pl. Sys. And Evo.*, Vol. 169, [Issue-3](#), pp 209-217.

Benoit Lafleur, Linda M. Hooper Bu, Paul Mumma, James P. Geaghan. 2005. Soil fertility and plant growth in soils from pine forests and plantations: Effect of invasive red imported fire ants *Solenopsis invicta* (Buren), *Pedobiologia* [Volume 49, Issue 5](#), 3 October Pg.415–423.

Cammeraat L H, Willott S J, Compton S G, Geoderma L D, 2002. The effects of ants' nests on the physical, chemical and hydrological properties of a rangeland soil in semi-arid Spain. *Geoderma*, Vol. 105, Issues 1–2, January 2002, Pages 1–20.

Diane Wagner, Mark J. F. Brown and Deborah M. Gordon, **1997.** Harvester ant nests, soil biota and soil chemistry. *Oecologia*, Vol. 112, [Issue 2](#), pp 232-236.

Diane Wagner and Jeremy B. Jones (2005). The Contribution of Harvester Ant Nests, *Pogonomyrmex rugosus* (Hymenoptera, Formicidae), to Soil Nutrient Stocks and Microbial Biomass in the Mojave Desert. *Oecologia*, Vol.2,pg223-232..

Domisch J. Finer L. Ohashi M. Risch A.C. Sundstrom L., Niermela P. and Jurgenser M. F., 2006. Contribution of Red wood ant mounds to forest floor CO₂ in boreal Coniferous soil.,- *Biology and Biochemistry*, 38: Pg. 2425-2433.

Eldridge, D.J. and Pickard, J. (1994). The effect of ants on sandy soils in semi-arid eastern Australia. II. Nest turnover and consequences for bioturbation. *Australian Journal of Soil Research* 32, 323-333.

Eldridge, D.J. and Myers, C.A. (1998). Some evidence that the ants *Aphaenogaster barbigula* (Myrmicinae) enhance soil nutrients around their nest entrances in semi-arid eastern Australia. *Australian Journal of Soil Research* 36, 1009-1017.

Folgarait P.J., 1998. Ant biodiversity and its relationship to ecosystem functioning a review. *Biodiv. and Conserv.*, 7: Pg. 1221-1224.

Frouz J. and Kalick J. 1996. The role of ants for availability of phosphorous in spruce plantation proceedings of the national conf. about study of phosphorous and other elements. *Pedobiologia*, 75-80.

Frouz J., Holec M. and Kalick J. 2003. The effect of *Lasius niger* (Hymenoptera: Formicidae) ant nest on related soil chemical properties. *Pedobiologia* 47: 205-212.

Kadu S.G. (2010). Biology and Bioinsecticidal Control of Carpenter Ant, *Camponotus compressus* (Hymenoptera-Formicidae) *Ph. D. Thesis*, RTM University, Nagpur, M.S., India.

D. D. Barsagade, D.B. Tembhare and Seema G. Kadu (2013). Microscopic structure of antennal sensilla in the carpenter ant *Camponotus compressus* (Fabricius) (Formicidae: Hymenoptera) (2013). *Asian Myrmecology*, Volume 5, 113–120.

Seema G. Kadu (2014). Study on Diversity and Species Richness of Ants In Nagpur Region (Ms), India. *IJBAT*, ISBN No. 2348-317, sp. Is. Dec. 2014.

Ohashi M, Kilpelainen, J., Finer L., Risch A.C., Domisch T. Neuvonen, S. and Niemela P. 2007a. The effect of red wood ant (*Formica rufa* group) mounds on roof biomass, density and nutrient concentration in boreal managed forests, *Jour., Forest research*, 12: Pg.113-119.