



---

## HIGHEST COMMON FACTOR BY THE VEDIC MATHEMATICS

**DR. SHARMILA KUMRI**  
**ASSISTANT PROFESSOR IN MATHEMATICS**  
**MAHILA MAHAVIDYALAYA JHOJHU KALAN**  
**CHARKHI DADRI (HARYANA)**  
[Kumarisharmila908@gmail.com](mailto:Kumarisharmila908@gmail.com)

**ABSTRACT :** This paper shows to calculate Highest common factors by Modern method and Vedic method lopna sthapanabhyam sutra sankalan - vyavakalabhyam process and the Adyamadya rule. It can also be applied in polynomial expressions.

### INTRODUCTION –

Finding Highest common factor of numbers and Algebraic expressions is fundamental process in Mathematics with much application. Highest common factor is also known as greatest common factor. There are two methods which are used to find HCF in Modern Mathematics. The first is by means of factorization and the second is by process of continuous division. The Vedic method provides a third method which is applicable to all cases.

**LOPNA-STHAPANABHYAM SUTRA** - It means ‘By Alternate elimination and retention or alternate destruction’ of the highest and the lowest powers.

### HCF BY MODERN METHODS

Take an example

Example 1: Find the H.C.F. of  $x^2 + 5x + 4$  and  $x^2 + 7x + 6$ .

1. Factorization method:

$$x^2 + 5x + 4 = (x + 4)(x + 1)$$

$$x^2 + 7x + 6 = (x + 6)(x + 1)$$

H.C.F. is  $(x + 1)$ .

2. Continuous division process.

$$\begin{array}{r}
 x^2 + 5x + 4 \quad x^2 + 7x + 6 \quad (1) \\
 \hline
 \phantom{x^2 + 5x + 4} \phantom{x^2 + 7x + 6} \phantom{(1)} x^2 + 5x + 4 \\
 \hline
 \phantom{x^2 + 5x + 4} \phantom{x^2 + 7x + 6} \phantom{(1)} \phantom{x^2 + 5x + 4} x^2 + x \\
 \hline
 \phantom{x^2 + 5x + 4} \phantom{x^2 + 7x + 6} \phantom{(1)} \phantom{x^2 + 5x + 4} \phantom{x^2 + x} 4x + 4 \quad 2x + 2 \quad (1/2) \\
 \hline
 \phantom{x^2 + 5x + 4} \phantom{x^2 + 7x + 6} \phantom{(1)} \phantom{x^2 + 5x + 4} \phantom{x^2 + x} \phantom{4x + 4} 2x + 2 \\
 \hline
 \phantom{x^2 + 5x + 4} \phantom{x^2 + 7x + 6} \phantom{(1)} \phantom{x^2 + 5x + 4} \phantom{x^2 + x} \phantom{4x + 4} \phantom{2x + 2} 0
 \end{array}$$

Thus  $4x + 4$  i.e.,  $(x + 1)$  is H.C.F.

### HIGHEST COMMON FACTOR BY VEDIC METHODS

Lopana - Sthapana process i.e. elimination and retention or alternate destruction of the highest and the lowest powers is as below:

$$\text{Subtract} \quad \left\{ \begin{array}{l} x^2 + 5x + 4 \\ x^2 + 7x + 6 \\ \hline -2) -2x - 2 \\ \hline x + 1 \end{array} \right.$$

i.e.,  $(x + 1)$  is the H.C.F.

Example 2: Find H.C.F. of  $2x^2 - x - 3$  and  $2x^2 + x - 6$

$$\text{Subtract} \quad \left\{ \begin{array}{l} 2x^2 - x - 3 \\ 2x^2 + x - 6 \\ \hline -1) -2x + 3 \\ \hline 2x + 3 \end{array} \right. \text{ is the H.C.F}$$

Example 3:  $x^3 - 7x - 6$  and  $x^3 + 8x^2 + 17x + 10$ .

Now by Lopana - Sthapana and Sankalana - Vyavakalanabhyam

$$\text{Subtract} \quad \left\{ \begin{array}{l} x^3 - 7x - 6 \\ x^3 + 8x^2 + 17x + 10 \\ \hline -8) -8x^2 - 24x - 16 \\ \hline x^2 + 3x + 2 \end{array} \right.$$

Example 4:  $x^3 + 6x^2 + 5x - 12$  and  $x^3 + 8x^2 + 19x + 12$ .

$$\text{Add} \quad \left\{ \begin{array}{l} x^3 + 6x^2 + 5x - 12 \\ x^3 + 8x^2 + 19x + 12 \\ \hline 2x^3 + 14x^2 + 24x \\ \hline 2x(x^2 + 7x + 12) \end{array} \right.$$

(or)

$$\begin{array}{r} \text{Subtract} \\ \left. \begin{array}{l} x^3 + 6x^2 + 6x - 12 \\ x^3 + 8x^2 + 19x + 12 \\ \hline -2x^2 - 14x - 24 \end{array} \right\} \\ -2x(x^2 + 7x + 12) \end{array}$$

So  $x^2 + 7x + 12$  is highest common factor

Now Algebraic Proof:

Let P and Q be two expressions and H is their H.C.F. Let A and B the Quotients after their division by H.C.F.

$$\text{i.e., } \frac{P}{H} = A \text{ and } \frac{Q}{H} = B \text{ which gives } P = A.H \text{ and } Q = B.H$$

$$P + Q = AH + BH \text{ and } P - Q = AH - BH$$

$$= (A + B).H \qquad \qquad = (A - B).H$$

Thus we can write  $P \pm Q = (A \pm B).H$

Similarly  $MP = M.AH$  and  $NQ = N.BH$  gives  $MP \pm NQ = H (MA \pm NB)$

This states that the H.C.F. of P and Q is also the H.C.F. of  $P \pm Q$  or  $MA \pm NB$ .

i.e. we have to select M and N in such a way that highest powers and lowest powers (or Independent terms) are removed and H.C.F appears as we have seen in the examples.

## CONCLUSION

Although Modern methods 1<sup>st</sup> is Factorisation method and 2<sup>nd</sup> is process of continuous division are more intellectual but harder to work as high power of algebraic into further factors is greater waste of time , energy and less dependable. The Vedic method is free from all these defects and is not only intellectual but also simple, easy and reliable .

## REFERENCES

- 1 . Bharti Krishan Tirthji Maharaj , **Vedic Mathematics** , Motilal Banarsidass publisher ,2012, page no. 87 – 91.
2. Kenneth Williams , “HIGHEST COMMON FACTOR AND LOWEST COMMON MULTIPLE USING THE SESA SUTRA” , **online International journal of Vedic Mathematics** , serial no. 53 , Vedic Mathematics Academy ,U.K. , 2021.
3. Sharmila kumari , “ Importance of Vedic Mathematics and its role in Mathematics” , Thesis (unpublished), Mathematics ,Singhania University ,Pacheri Berri, 2015.